



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OPP OFFICIAL RECORD
HEALTH EFFECTS DIVISION
SCIENTIFIC DATA REVIEWS
EPA SERIES 361

OFFICE OF
PREVENTION, PESTICIDES, AND
TOXIC SUBSTANCES

TXR No. 0054792

MEMORANDUM

DATE: December 12, 2007

SUBJECT: **Thiencarbazone-Methyl:** Qualitative Risk Assessment Based On
C57BL/6J Mouse Carcinogenicity Dietary Study

P.C. Code: 015804

TO: John Doherty, Toxicologist
Reregistration Branch 3
Health Effects Division (7509P)

FROM: Lori L. Brunsman, Statistician
Science Information Management Branch
Health Effects Division (7509P)

THROUGH: Jessica Kidwell, EPS
and
Jess Rowland, Branch Chief
Science Information Management Branch
Health Effects Division (7509P)

BACKGROUND

A carcinogenicity study in C57BL/6J mice was conducted by Bayer CropScience, Sophia Antipolis Cedex, France, for Bayer AG, Bayer CropScience, Monheim, Germany, and completed November 10, 2006 (Study No. SA 04062, MRID No. 47070135).

The study design allocated groups of 50 mice to dose levels of 0, 200, 1000 and 4000 ppm of Thiencarbazone-methyl for 78 weeks. Doses were equivalent to 0, 29.2, 147 and 599 mg/kg/day for males and 0, 36.8, 185 and 758 mg/kg/day for females. An additional ten mice per sex per dose were designated for interim sacrifice at week 29.

DEC 20 REC'D 2007
EHC
SH

ANALYSES

Survival Analyses

Male mice showed a statistically significant increasing trend for mortality with increasing doses of Thiencarbazone-methyl, as well as a significant pair-wise comparison of the 4000 ppm dose group with the controls, both at $p < 0.05$. There was not a statistically significant trend in mortality with increasing doses of Thiencarbazone-methyl in female mice, however, there was a statistically significant pair-wise comparison of the 200 ppm dose group with the controls at $p < 0.05$ (Tables 1 and 2).

Tumor Analyses

Male mice had statistically significant trends in bladder transitional cell papillomas and carcinomas, both at $p < 0.05$. There was also a statistically significant trend at $p < 0.01$, and a significant pair-wise comparison of the 4000 ppm dose group with the controls at $p < 0.05$, for bladder transitional cell papillomas and carcinomas combined. Female mice had a statistically significant trend in bladder transitional cell papillomas and carcinomas combined at $p < 0.05$, however, female mice had no significant pair-wise comparisons of the dosed groups with the controls. The statistical analyses of the tumors in the male mice were based upon Peto's Prevalence Test. The statistical analyses of the tumors in the female mice were based upon Fisher's Exact Test for pair-wise comparisons and the Exact Test for trend (Tables 3 and 4).

Table 1. Thien carbazon-Methyl – C57BL/6J Mouse Study (MRID 47070135)

Male Mortality Rates⁺ and Cox or Generalized K/W Test Results

Dose (ppm)	<u>Weeks</u>				Total
	1-28	29 ⁱ	29-53	54-80 ^f	
0	0/60	10/60	4/50	7/46	11/50 (22)*
200	4/60	9/56	1/47	12/46	17/51 (33)
1000	1/60	9/59	3/50	8/47	12/51 (24)
4000	1/60	10/59	6/49	15/43	22/50 (44)*

⁺Number of animals that died during interval/Number of animals alive at the beginning of the interval.

ⁱInterim sacrifice at week 29.

^fFinal sacrifice at weeks 78-80.

()Percent.

Note: Time intervals were selected for display purposes only.
 Significance of trend denoted at control.
 Significance of pair-wise comparison with control denoted at dose level.
 If *, then $p < 0.05$. If **, then $p < 0.01$.

Table 2. Thiencarbazon-Methyl – C57BL/6J Mouse Study (MRID 47070135)

Female Mortality Rates⁺ and Cox or Generalized K/W Test Results

Dose (ppm)	<u>Weeks</u>				Total
	1-28	29 ⁱ	29-53	54-80 ^f	
0	0/60	10/60	3/50	3/47	6/50 (12)
200	2/60	10/58	2/48	10/46	14/50 (28)*
1000	1/60	10/59	1/49	2/48	4/50 (8)
4000	0/60	10/60	1/50	8/49	9/50 (18)

⁺Number of animals that died during interval/Number of animals alive at the beginning of the interval.

ⁱInterim sacrifice at week 29.

^fFinal sacrifice at weeks 78-80.

()Percent.

Note: Time intervals were selected for display purposes only.
 Significance of trend denoted at control.
 Significance of pair-wise comparison with control denoted at dose level.
 If *, then $p < 0.05$. If **, then $p < 0.01$.

Table 3. Thiencarbazone-Methyl – C57BL/6J Mouse Study (MRID 47070135)

Male Urinary Bladder and Urethra/Prostate Transitional Cell Tumor Rates⁺
and Peto's Prevalence Test Results

	Dose (ppm)			
	0	200	1000	4000
Bladder Papillomas (%)	0/39 (0)	0/34 (0)	0/39 (0)	1 ^a /28 (4)
p =	0.02686*	-	-	0.11896
Urethra Carcinomas (%)	0/38 (0)	0/34 (0)	0/37 (0)	1 ^b /28 (4)
p =	0.02835*	-	-	0.12202
Combined (%)	0/38 (0)	0/34 (0)	0/37 (0)	2/28 (7)
p =	0.00342**	-	-	0.04842*

+Number of tumor bearing animals/Number of animals examined, excluding those that died or were sacrificed before observation of the first tumor.

^aFirst urinary bladder transitional cell papilloma observed at week 80, dose 4000 ppm, in a final sacrifice animal.

^bFirst urethra/prostate transitional cell carcinoma observed at week 79, dose 4000 ppm, in a final sacrifice animal.

Note: Significance of trend denoted at control.
Significance of pair-wise comparison with control denoted at dose level.
If *, then $p < 0.05$. If **, then $p < 0.01$.

Table 4. Thiencarbazon-Methyl – C57BL/6J Mouse Study (MRID 47070135)

Female Bladder Transitional Cell Tumor Rates⁺
and Fisher's Exact Test and Exact Test for Trend Results

	Dose (ppm)			
	0	200	1000	4000
Papillomas (%)	0/45 (0)	0/45 (0)	0/46 (0)	2 ^a /48 (4)
p =	0.06700	1.00000	1.00000	0.26367
Carcinomas (%)	0/45 (0)	0/45 (0)	0/46 (0)	1 ^b /48 (2)
p =	0.2609	1.00000	1.00000	0.51613
Combined (%)	0/45 (0)	0/45 (0)	0/46 (0)	3/48 (6)
p =	0.01693*	1.00000	1.00000	0.13329

+Number of tumor bearing animals/Number of animals examined, excluding those that died or were sacrificed before week 54.

^aFirst papilloma observed at week 79, dose 4000 ppm, in a final sacrifice animal.

^bFirst carcinoma observed at week 79, dose 4000 ppm, in a final sacrifice animal.

Note: Significance of trend denoted at control.
 Significance of pair-wise comparison with control denoted at dose level.
 If *, then p < 0.05. If **, then p < 0.01.

References

- Cox, D.R. (1972) Regression Models and Life Tables (with discussion). J. Royal Stat. Soc. Ser. B. 34, 187-220.
- Gart, J.J., D. Krewski, P.N. Lee, R.E. Tarone, and J. Wahrendorf (1986) The Design and Analysis of Long-Term Animal Experiments. In: Statistical Methods in Cancer Research, Volume III. IARC Scientific Publications No. 79. Lyon, France: International Agency for Research on Cancer, p. 18.
- Peto, R., M. Pike, N. Day, R. Gray, P. Lee, S. Parish, J. Peto, S. Richard, and J. Wahrendorf (1980) Guidelines for Simple, Sensitive, Significant Tests for Carcinogenic Effects in Long-Term Animal Experiments. In: Monographs on the long-term and short-term screening assays for carcinogens: a critical appraisal. IARC Monographs, Supplement 2. Lyon, France: International Agency for Research on Cancer, pp. 311-426.
- Thomas, D.G., N. Breslow, and J.J. Gart (1977) Trend and Homogeneity Analyses of Proportions and Life Table Data. Computers and Biomedical Research 10, 373-381.



13544



R155878

Chemical: Thien carbazone-methyl

PC Code:
015804

HED File Code: 31000 Qualitative Analysis Reviews

Memo Date: 12/12/2007

File ID: TX0054792

Accession #: 000-00-0124

HED Records Reference Center
1/15/2008

